



**Focused on  
Growth and Innovation**

**“Patients are at the heart  
of what we do”**

Investor presentation  
February 24, 2021



# Table of contents

- Executive summary
- Five reasons to invest
- Portfolio
  - Antifungal
    - Cresemba® (isavuconazole)
  - Antibiotic
    - Zevtera® (ceftobiprole)
  - Oncology
    - Derazantinib
    - Lisavanbulin (BAL101553)
- Financials
- Appendix





## Executive summary



# Experienced leadership team



**David Veitch** CEO

Joined 2014

Previous roles:



**Adesh Kaul** CFO

2009



**Marc Engelhardt** MD, Ph.D. CMO

2010



**Gerrit Hauck** Ph.D. CTO

2018



**Laurenz Kellenberger** Ph.D. CSO

2000




# At a glance

- Well funded, commercial-stage biopharmaceutical company with significantly growing cash flows from commercialized products
- Focused in the areas of oncology and infectious diseases
- Potential for sustainable growth and value creation based on commercialized brands and an innovative pipeline
- Experienced people with the proven expertise to take compounds from research to market
- Two revenue generating hospital anti-infective brands, Cresemba® and Zevtera® and two clinical oncology drug candidates
- Recognized ability to establish and manage partnerships in both the development and commercial phase, providing access to international markets
- Listed on SIX Swiss Stock Exchange, SIX: BSLN
- Based in life sciences hub, Basel, Switzerland



# Potential for sustainable growth and value creation based on commercialized brands and innovative pipeline




# Our strategy



## Foster

Foster an agile organisation based on a dynamic and open culture



## Focus

Focus on continuously increasing cash flow from our two commercial-stage hospital anti-infective brands, Cresemba<sup>®</sup> and Zevtera<sup>®</sup>



## Leverage

Leverage our expertise in bringing drugs from research to market by utilising appropriate partnerships with established organisations



## Invest


Invest in our clinical portfolio of targeted, small molecule, oncology drug candidates and the phase 3 ceftobiprole program



## Innovate

Continue to broaden our R&D pipeline through both internal and external innovation

# Global coverage — Cresemba®





# The company we keep — established strong partnerships

## License partners



Europe (excl. Nordics), China  
Asia-Pacific, Russia, Turkey  
and Israel (Cresemba®)



U.S. (Cresemba®)



Japan (Cresemba®)



China (Zevtera®)

## Distribution partners



Europe (excl. Nordics), Israel  
(Zevtera®)



MENA region  
(Cresemba® and Zevtera®)




LatAm  
(Cresemba® and Zevtera®)



Nordics  
(Cresemba® and Zevtera®)



Canada  
(Cresemba® and Zevtera®)





**Five reasons  
to invest**



# Five reasons to invest



## Growth

Well funded with increasing and sustainable cash flow through commercialized brands



## Prospects

Opportunity to share in pipeline value creation and proven approach to the successful commercialization of products around the world



## Leadership

Experienced team working in an agile culture able to turn pipeline projects into revenue generating brands



## Partnerships

Proven ability to build successful partnerships in research, development and commercialization with leading academic, governmental and industrial organisations



## Focus

One of the few biopharmaceutical companies in the world focused on the development and commercialization of targeted oncology small molecules and new antibiotics and antifungals

Antifungal

**Cresemba<sup>®</sup>**  
**(isavuconazole)**


Invasive mold infections



# The market — Invasive fungal infections


- Severe, potentially life-threatening infections mainly affecting immunocompromised patients
- An important cause of morbidity and mortality in cancer patients undergoing intensive chemotherapy regimens
- Rising number of immunocompromised patients (cancer and transplantations) driving therapeutic demand
- Mucorales infections on the rise – doubled from 2000 to 2013
- Limitations of current therapies (spectrum of activity, toxicity, effective plasma levels) drive the need for new agents

## Mortality rates for invasive fungal infections\*\*



\*\*Kullberg/Arendrup *N Engl J Med* 2015, Baddley *Clin Infect Dis* 2010, Roden *Clin Infect Dis* 2005, Greenberg *Curr Opin Infect Dis* 2004

# Cresemba continues strong in-market sales uptake



USD 244 mn  
 “in-market”  
 sales in the  
 12 months to  
 September  
 2020


LCD: USD corrected for currency fluctuations; MAT: Moving annual total; Source: IQVIA, September 2020

# Sales of best-in-class antifungals\* by product

USD 3.0 bn sales (MAT Q3 2020)

- Potential to increase Cresemba® (isavuconazole) market share
  - Anticipate to be launched in 60 countries by end-2021
  - Exclusivity through 2027 in the U.S. and potential pediatric exclusivity extension to 2027 (from 2025) in the EU

\* Best-in-class antifungals: isavuconazole, posaconazole, voriconazole, AmBisome, anidulafungin, caspofungin, micafungin



MAT: Moving annual total; Sales figures in USD, corrected for currency fluctuations; Source: IQVIA, September 2020

Confidential/proprietary information of Basilea Pharmaceutica International Ltd. – not for distribution

# Cresemba<sup>®</sup> — Differentiated by spectrum, safety and tolerability

- Broad spectrum of activity against molds, including emerging molds (mucorales)
- Consistent plasma levels
- Statistically fewer drug-related adverse events and treatment-emergent adverse events (liver, skin, eye) in invasive aspergillosis patients vs. voriconazole in SECURE phase 3 study
- Can be administered without restriction in patients with renal impairment
- Manageable drug-drug interaction profile
- Once daily maintenance dose, i.v./oral treatment
- ECIL-6 guideline: Cresemba<sup>®</sup> recommended for the first-line treatment of invasive aspergillosis in leukemia and hematopoietic stem cell transplant patients. ECIL states that isavuconazole is as effective as voriconazole with a better safety profile.



Antibacterial  
**Zevtera<sup>®</sup>**  
**(ceftobiprole)**

Severe bacterial infections



# Zevtera<sup>®</sup> — An introduction

- Broad-spectrum anti-MRSA cephalosporin (including Gram-negative bacteria)
- Rapid bactericidal activity
- Potential to replace antibiotic combinations
- Early improvement in HAP, particularly in patients with MRSA, and CAP, including high-risk patients
- Cephalosporin class safety profile
- Marketed in selected countries in Europe, Latin America and the MENA-region as well as in Canada


Approved in major European countries & several non-European countries for both hospital-acquired pneumonia (HAP), excluding ventilator-associated pneumonia (VAP), and community-acquired pneumonia (CAP). Not approved in the U.S.

MENA: Middle East and North Africa




# The hospital anti-MRSA antibiotic market — A USD 2.8 bn market\* with the U.S. being the most important region

Daptomycin sales by region (2015, before LOE)



Ceftaroline sales by region (MAT Q3 2020)



\* Vancomycin, linezolid, teicoplanin, daptomycin, tigecycline, telavancin, ceftaroline, dalbavancin, oritavancin, and tedizolid

MRSA: Methicillin-resistant *Staphylococcus aureus*; LOE: Loss of exclusivity; ROW: Rest of world  
MAT: Moving annual total; Sales figures in USD, corrected for currency fluctuations; Source: IQVIA, September 2020

# Strategy for accessing the U.S. market

- Two cross-supportive phase 3 studies under FDA Special Protocol Assessment (SPA)
- Phase 3 program largely funded by BARDA (up to USD ~130 mn, ~70% of total program costs)

1. Acute Bacterial Skin and Skin Structure Infections (ABSSSI)<sup>1</sup> successfully completed



2. *Staphylococcus aureus* bacteremia (SAB)<sup>2</sup> ongoing, topline results from phase 3 study expected in H1 2022



- Qualified Infectious Disease Product (QIDP) designation extends U.S. market exclusivity to 10 years from approval

<sup>1</sup> Overcash JS et al. ECCMID 2020, abstract 1594. (NCT03137173)

<sup>2</sup> Hamed K et al. Future Microbiol. 2020;15:35-48. (NCT03138733)

# SAB – an area with high medical need

- Nearly 120,000 *S. aureus* bloodstream infections in the US (in 2017)<sup>1</sup>
- ERADICATE targets complicated SAB, characterized by concomitant or metastatic infections such as bone, joint or heart valve infections; persistent bacteremia; or bacteremia in patients on dialysis
- Substantial morbidity and approximately 20% 30-day mortality<sup>2</sup>
- Limited antibiotic treatment options with only two approved treatments for SAB in the U.S. that cover both MSSA and MRSA, i.e. vancomycin and daptomycin


<sup>1</sup> MMWR, 2019;68:214–219.

<sup>2</sup> Hamed K et al. Future Microbiol. 2020;15:35-48.

MRSA: methicillin-resistant *Staphylococcus aureus*

MSSA: methicillin-susceptible *Staphylococcus aureus*

## Causes and consequences of SAB



Adapted from Edwards AM et al. Trends Microbiol. 2011;19:184-190.

A microscopic image of cells, likely cancer cells, with an orange overlay. The cells are spherical and have a textured surface. Some cells are larger and more prominent than others. The background is a dense network of fine, fibrous structures. The overall color scheme is dominated by shades of orange and yellow.


Oncology

# Derazantinib

FGFR-driven tumors

# Targeting FGFR-driven tumors as single agent and in combination with immunotherapy

- Small molecule, oral inhibitor of FGFR family of kinases
- Development strategy focused on achieving differentiation by leveraging unique properties of derazantinib
  - Kinase inhibition profile: exploring therapeutic potential of additional targets of derazantinib such as CSF1R and VEGFR2 kinase
  - Safety profile: exploring relevance for potential combination therapies
- Three clinical studies ongoing
  - FIDES-01 (Ph 2) in intrahepatic cholangiocarcinoma (iCCA)
  - FIDES-02 (Ph 1/2) in urothelial cancer
  - FIDES-03 (Ph 1/2) in gastric cancer



Sources: Helsten et al., Clin Cancer Res 2016 (22), 257-267; FGFR2 fusions in iCCA: Graham et al. Hum Pathol 2014 (45), 1630-1638; Jain et al. JCO Precis Oncol 2018 (2) 1-12

# Phase 2 study in iCCA\* – established clinical proof of concept in FIDES-01

FIDES-01 Cohort 1 (N=103)
<p><b>FGFR2 fusions</b> (~15% of iCCA)</p> <p><b>Topline results:</b> ORR: 20.4% Median PFS: 6.6 months</p>

- Consistent with earlier Phase 1/2 data<sup>1</sup> and with interim analysis from FIDES-01
- Clinical proof of concept for derazantinib as monotherapy in its first indication established

FIDES-01 Cohort 2 (N=43) - ongoing
<p><b>FGFR2 mutations/amplifications</b> (~5% of iCCA)</p> <p>Pooled data from 23 patients (clinical studies/EAP)<sup>2</sup> Median PFS 7.2 months</p>

- Encouraging progression-free survival in pooled analysis consistent with outcome in patients with FGFR2 gene fusions
- Interim results expected H1 2021

**Manageable safety profile with low incidence of nail toxicity, retinal events, hand-foot syndrome and stomatitis**

FIDES-01: NCT03230318

<sup>1</sup>Mazzaferro et al. Br J Cancer. 2019

<sup>2</sup>Droz Dit Busset et al. Annals of Oncology (2020) 31 (suppl\_5): abstract 45P (NCT01752920, NCT03230318)

\*in patients who progressed after at least one prior systemic chemotherapy regimen



# Clinical program in urothelial cancer – FIDES-02

Multi-cohort phase 1b/2 study of derazantinib monotherapy or in combination with atezolizumab in patients with urothelial cancer harboring FGFR genetic aberrations

- Substudies (N≈300) in various treatment settings, including:
  - Post-chemotherapy/immunotherapy recurrence (second-line and post second-line)
  - First-line platinum-ineligible, PD-L1-low
  - Resistance to prior FGFR-inhibitor treatment
- Successful completion of phase 1b cohort
  - Recommended phase 2 dose for the combination established
  - No dose-limiting toxicities observed
- Clinical supply agreement with Roche for atezolizumab
- Plan to amend the study protocol to explore a higher dose of derazantinib in two cohorts of this study
  - Supported by the observed safety and tolerability profile of derazantinib at the current dose of 300 mg per day
  - May provide additional benefits in monotherapy and combination to patients with FGFR-positive urothelial cancer
  - Considers the evolving treatment and competitive landscape in urothelial cancer in patients both with and without FGFR genetic aberrations
- Interim results in derazantinib monotherapy expected H1 2021
- Interim results in combination therapy with atezolizumab expected H2 2021

# Clinical program in gastric cancer – FIDES-03

Multi-cohort Phase 1b/2 study of derazantinib as monotherapy or in combination therapy with standard of care (ramucirumab/paclitaxel) or atezolizumab in patients with advanced HER2-negative gastric adenocarcinoma harboring FGFR genetic aberrations

- Substudies using derazantinib monotherapy or combination treatment, including:
  - Derazantinib monotherapy in various molecular subtypes
  - Combination of derazantinib with ramucirumab/paclitaxel
  - Combination of derazantinib with atezolizumab
- Clinical supply agreement with Roche for atezolizumab
- Clinical trial collaboration and supply agreement with Lilly for ramucirumab
- Interim results in derazantinib monotherapy and recommended phase 2 dose of derazantinib in combination with ramucirumab/paclitaxel expected H2 2021

# FGFR-inhibitors show differences in safety profiles

	Cholangiocarcinoma				Urothelial cancer	
	DZB <sup>1</sup> (N=44)	INF <sup>2</sup> (N=71)	FUT <sup>3</sup> (N=67)	PEM <sup>4</sup> (N=146)	PEM <sup>5</sup> (N=108)	ERD <sup>6</sup> (N=87)
Dosing regimen	300mg QD	125mg Q4W QD for 3w	20 mg QD	13.5mg Q3W QD for 2w	13.5mg Q3W QD for 2w	8 mg QD (titration to 9mg)
Most frequent safety events	Phosphorus↑ Nausea Vomiting	Phosphorus↑ Fatigue Stomatitis	Phosphorus*↑ Diarrhea* Dry mouth*	Phosphorus↑ Alopecia Diarrhoea	Diarrhoea Alopecia Constipation	Phosphorus↑ Stomatitis Fatigue
Blood phosphorus↑†	59%	73%	88%	60%	31%	76%
Fatigue†	43%	49%	NR	42%	32%	54%#
Alopecia†	20%	38%	NR	49%	40%	26%
Dry eye/xerophthalmia†	16%	32%	NR	35%#	NR	28%#
Retinopathy†	0%	NR	9%	6%‡	NR	25%
Alanine aminotransferase (ALT) ↑	30%**	NR	NR	43%**	NR	41%**
Hand-foot syndrome/PPE	0%	27%	18%	15%	NR	26%
Nail toxicities	<5%	NR	42%	43%#	NR	41%#
Stomatitis	11%	45%	NR	35%	34%	56%

<sup>1</sup> Droz Dit Busset et al., ESMO 2019 and Basilea data on file, <sup>2</sup> Javle et al., ESMO 2018, <sup>3</sup> Goyal et al., ASCO 2020, <sup>4</sup> Pemazyre™ U.S. Prescribing Information (April 2020), <sup>5</sup> Necchi, et al., ESMO 2018,

<sup>6</sup> Balversa™ U.S. prescribing information (April 2019)

† assumed FGFR inhibitor class-effect; \*futibatinib treatment-related adverse events

# includes various and different adverse reactions; for details see Pemazyre™ U.S. Prescribing Information (April 2020) and Balversa™ U.S. prescribing information (April 2019);

† Refers to reported adverse events of Retinal Pigment Epithelial Detachment (RPED) for pemigatinib, Central Serous Retinopathy (CSR)/RPED for erdafitinib and CSR for futibatinib

‡ reported incidence is from 466 patients who received Pemazyre™ across clinical trials;

\*\* based on reported adverse events for DZB; based on reported laboratory abnormalities, regardless of causality for PEM and ERD.

Abbreviations: DZB: derazantinib, INF: infigratinib (BGJ398), FUT: futibatinib (TAS-120), PEM: pemigatinib (INCB54828), ERD: erdafitinib; PPES: Palmar-plantar erythrodysesthesia; NR: not reported; QD: daily; Q3W/Q4W: every 3/4 weeks; w: weeks

Oncology


# Lisavanbulin (BAL101553)

Glioblastoma  
and other solid tumors



# Novel tumor checkpoint controller crossing the blood-brain barrier

- Novel compound inducing tumor cell death through spindle assembly checkpoint activation
- Targeting diverse tumor types resistant to standard therapeutic approaches
- Flexible dosing potential, including daily oral dosing
- Comprehensive biomarker program to optimize patient selection
- Crosses the blood-brain barrier with potent activity in brain tumor models alone and in combination



\* Lisavanbulin (BAL101553) is a prodrug of BAL27862


Confidential/proprietary information of Basilea Pharmaceutica International Ltd. – not for distribution

# Biomarker-driven phase 2 study ongoing in recurrent glioblastoma


- EB1 (end-binding protein 1) is located on the microtubules and involved in microtubule dynamics and has been shown to be a response predictive marker for lisavanbulin in preclinical studies
- Strong EB1 staining was observed in a patient with an exceptional response to daily oral lisavanbulin in the phase 1 dose-escalation study in recurrent glioblastoma<sup>1</sup>
  - Patient ongoing for more than two years
  - >80% reduction in glioblastoma tumor size
- Interim results expected H2 2021

<sup>1</sup> Lopez et al. Phase 1/2a study of once daily oral BAL101553, a novel tumor checkpoint controller, in adult patients with progressive or recurrent glioblastoma or high-grade glioma. JCO 2019;37:15 suppl, 2025 (NCT02490800)


## Glioblastoma tumor size reduction in an exceptional responder and EB1 staining of glioblastoma tissue compared to non-responding patients




Baseline (May 2018)



Post Cycle 12 (April 2019)



Responder




Non-responder


# Unmet medical need in glioblastoma

- The most common primary brain cancer in adults with an incidence of 3-4 per 100,000 people, (though geographic variation exists) and a median age at onset of > 60 years
- Associated with poor prognosis, high morbidity and healthcare burden
- 5-year survival is below 5% with current standard of care (multimodality treatment including surgery, radiotherapy, chemotherapy)<sup>1</sup>
- *MGMT*-promoter methylation status has been demonstrated as a predictor for the response to (radio)chemotherapy (temozolomide)<sup>2</sup>
- Established molecular markers used for classification include *IDH* mutations and/or 1p/19q codeletion<sup>3</sup>
- No molecular targeted therapy currently approved

## Radiological and tissue presentations of glioblastoma



Variable glioblastoma appearances on post-gadolinium T1-weighted MRI: central necrotic mass with nodular rim enhancement (a,b), predominantly solid enhancement (c), lack of contrast uptake (d)



Histological glioblastoma; H&E stain.

100 μm

Histological features of glioblastoma include marked hypercellularity, nuclear atypia, microvascular proliferation, and necrosis (N: necrosis, VP: vascular proliferations, T: tumor)

<sup>1</sup>Poon MTC et al. 2020; Sci Rep 10, 11622; <sup>2</sup>Hegi et al. NEJM 2005;352:997-1003

<sup>3</sup>Louis DN et al. Acta Neuropathol. 2016;131:803-820




# Financials








# Financial summary, in CHF mn (1/2)



Note: Consolidated figures in conformity with U.S. GAAP; rounding applied consistently

# Financial summary, in CHF mn (2/2)



Note: Consolidated figures in conformity with U.S. GAAP; rounding applied consistently

\*Cash, cash equivalents, restricted cash and investments


# Divestment of Chinese R&D subsidiary to U.S.-based custom manufacturing organization PHT International Inc. (“PHT”)

- Total purchase price of USD 6.3 million
  - USD 2.5 million upon closing, USD 3.8 million over the course of the next three years
- Basilea has entered into an agreement with PHT for continued R&D service provision
  - All 72 employees and the facilities will be transferred to PHT
  - Ensuring continuity on R&D projects
  - Providing sufficient time to optimize external sourcing of R&D services
- Closing of the transaction is expected in Q2 2021
- Financial impact
  - Annual operating expenses related to Chinese subsidiary mid-single digit million
  - Small positive P/L impact on sale in 2021
  - Limited (positive) impact in 2021/2022 on operating expenses due to transition period
  - Subsequently a greater potential impact on operating expenses

# Financial guidance

In CHF mn	FY 2020	FY 2021 guidance
Total revenue	127.6	128 - 138
thereof: Contributions Cresemba® & Zevtera® non-deferred	78.2	108 - 118
deferred	33.8	2.5
Operating loss	8.2	13 - 23
Cash and investments*	167.3	155 – 160**

Continued strong double-digit growth in Cresemba & Zevtera non-deferred revenue contributions Y-o-Y, CHF mn



\*Cash, cash equivalents, restricted cash and investments / \*\*Excluding any potential impact from a reduction of the outstanding convertible bonds

# Milestones & Outlook 2021 / 2022

**Cresemba® & Zevtera® — Increasing cash flows**  
**By the end of 2021, Cresemba to be on the market in 60 countries**


		H1 2021	H2 2021	H1 2022	H2 2022
<b>Isavuconazole</b>		✓ Complete patient enrolment in phase 3 study in Japan	Topline results from phase 3 study in Japan		
<b>Ceftobiprole</b>			Complete patient enrolment in SAB phase 3 study	Topline results from SAB phase 3 study	
<b>Derazantinib</b>	<b>FIDES-01 (iCCA)</b>	✓ Topline results (FGFR2 fusions)			
		Interim results (other FGFR2 gene aberrations)		Topline results (other FGFR2 gene aberrations)	
	<b>FIDES-02 (urothelial cancer)</b>	Interim results in derazantinib monotherapy	Interim results in combination therapy with atezolizumab		Topline results in combination therapy with atezolizumab
	<b>FIDES-03 (gastric cancer)</b>		Interim results in monotherapy and recommended phase 2 dose with ramucirumab/paclitaxel		Interim results in combination with ramucirumab/paclitaxel
<b>Lisavanbulin</b>			Interim results from phase 2 biomarker-driven glioblastoma study	Topline results from phase 2 biomarker-driven glioblastoma study	
			Recommended phase 2 dose in phase 1 study in newly-diagnosed glioblastoma in combination with radiotherapy		

# Appendix

# Significant sales of best-in-class antifungals in all major regions — Covered by our partnerships

USD 3.0 bn sales of best-in-class antifungals\* (MAT Q3 2020)

\* Best-in-class antifungals: isavuconazole, posaconazole, voriconazole, AmBisome, anidulafungin, caspofungin, micafungin



MAT: Moving annual total; Sales figures in USD, corrected for currency fluctuations; Source: IQVIA, September 2020

Confidential/proprietary information of Basilea Pharmaceutica International Ltd. – not for distribution


# Ceftobiprole — Positive topline phase 3 results reported in ABSSSI

Key topline study<sup>1</sup> results showing non-inferiority of ceftobiprole to vancomycin plus aztreonam for the primary and secondary endpoints



<sup>1</sup> NCT03137173  
ABSSSI: Acute bacterial skin and skin structure infections

## Early clinical response at 48–72h after start of treatment (ITT population)



ITT: intent-to-treat  
Pre-defined limit of non-inferiority = lower limit of 95 % CI for difference > -10 %



# Ceftobiprole — Positive topline phase 3 results reported in ABSSSI

Key topline study<sup>1</sup> results showing non-inferiority of ceftobiprole to vancomycin plus aztreonam for the primary and secondary endpoints




<sup>1</sup> NCT03137173

ABSSSI: Acute bacterial skin and skin structure infections

## Investigator-assessed clinical success at test-of-cure (TOC) 15-22 days after randomization (ITT, CE populations)

Patients with clinical success at the TOC visit (%)



CE: clinically evaluable; ITT: intent-to-treat

# Ceftobiprole key attributes for SAB treatment

- Beta-lactam antibiotic with rapid bactericidal activity against MSSA and MRSA<sup>1</sup>
- Superior activity profile in preclinical models of endocarditis compared to vancomycin and daptomycin<sup>2</sup>
- Low propensity for resistance development<sup>1</sup>
- Gram-negative coverage<sup>1</sup> in cases with polymicrobial infections
- Efficacy demonstrated in Phase 3 clinical trials in pneumonia and complicated skin and soft tissue infections<sup>1,3,4</sup>
- Established safety profile consistent with the cephalosporin class<sup>1,3</sup>


<sup>1</sup>Syed YY. *Drugs*. 2014;74:1523-1542.

<sup>2</sup>Tattevin P et al. *Antimicrob Agents Chemother*. 2010;54:610-613.

<sup>3</sup>Giacobbe DR et al. *Expert Rev Anti Infect Ther*. 2019;17:689-698.

<sup>4</sup>Overcash JS et al. *ECCMID 2020*, abstract 1594

## Comparative efficacy in a rabbit model of endocarditis



Organism titers in cardiac vegetations, spleens and kidneys of untreated and antibiotic treated rabbits infected with MRSA<sup>2</sup>

# Phase 3 study with ceftobiprole in the treatment of patients with SAB



- **Design:** randomized, double-blind, multi-center
- **Enrolment:** approximately 390 adult patients (male and female)
- **Indications:** *Staphylococcus aureus* bacteremia (SAB), including endocarditis (IE) and other forms of complicated SAB
- **Main inclusion criteria:** Positive *S. aureus* blood culture and signs & symptoms for SAB
- **Intervention:** ceftobiprole medocaril i.v.; comparator daptomycin i.v. or daptomycin plus aztreonam to cover Gram-negative bacteria
- **Primary endpoint:** overall success as assessed by an independent Data Review Committee (DRC) in the treatment of SAB, including IE, at the post-treatment evaluation (PTE) visit (70 days after randomization) in the modified intent-to-treat (mITT) population.
- **Secondary endpoints:** includes all-cause mortality at Day 28 and Day 70 (PTE visit) in the intent-to-treat (ITT) and mITT populations; and time to *S. aureus* bloodstream clearance


# FGFR-inhibitors show differences in kinase-inhibition profiles<sup>1</sup>

FGFR-inhibitor compound (Sponsor)	Parameter	FGFR1	FGFR2	FGFR3	FGFR4	CSF1R	VEGFR2
<b>Derazantinib</b> (Basilea)	Ratio to FGFR2 activity	4	1	4	77	3	6
<b>Pemigatinib</b> (Incyte)	Ratio to FGFR2 activity	3	1	4	39	231	62
<b>Erdafitinib</b> (Janssen)	Ratio to FGFR2 activity	2	1	2	13	95	6
<b>Rogaratinib</b> (Bayer)	Ratio to FGFR2 activity	5	1	6	18	116	48
<b>Infigratinib</b> (QED)	Ratio to FGFR2 activity	2	1	2	47	86	55
<b>Futibatinib</b> (Taiho)	Ratio to FGFR2 activity	2	1	2	18	NA	NA


<sup>1</sup> McSheehy et al. Derazantinib (DZB): A dual FGFR/CSF1R-inhibitor active in PDX-models of urothelial cancer. Mol Cancer Ther. 2019 (18) (12 Supplement) LB-C12

# In-silico analysis of derazantinib binding to CSF1R

- Crystal structures indicate differences in inhibitor binding sites of FGFR and CSF1R kinases
- Improved kinase inhibition activity of derazantinib against CSF1R versus other FGFR-inhibitors can be explained by the unique chemical structure of derazantinib<sup>1</sup>



**Derazantinib** (yellow) fits to smaller active site pocket of CSF1R (grey/red)




**Erdafitinib** (green) is too large (white circles) for the active site pocket of CSF1R (grey/red)

<sup>1</sup> McSheehy et al. Derazantinib (DZB): A dual FGFR/CSF1R-inhibitor active in PDX-models of urothelial cancer. Mol Cancer Ther. 2019 (18) (12 Supplement) LB-C12

# Potential therapeutic relevance of CSF1R-inhibition

- Derazantinib is active in inhibiting FGFR kinases and CSF1R (Colony-stimulating factor-1 receptor)
- CSF1R-inhibition may reprogram immunosuppressive tumor-infiltrating macrophages, restore T-cell activity and thereby improve the susceptibility to PD1/PD-L1 inhibitors<sup>1</sup>
- Derazantinib may address several oncogenic mechanisms at the same time, i.e. inhibiting FGFR and making the tumor more susceptible to immunotherapy
- Basilea entered into a clinical supply agreement with Roche to study a combination of derazantinib and Roche's PD-L1-blocking immune-checkpoint inhibitor atezolizumab in patients with urothelial and gastric cancer



<sup>1</sup> X. Zheng et al. Redirecting tumor-associated macrophages to become tumoricidal effectors as a novel strategy for cancer therapy. *Oncotarget*. 2017;8(29):48436-48452


Graph adapted from: A. Ghasemzadeh et al. New Strategies in Bladder Cancer: A Second Coming for Immunotherapy. *Clin Cancer Res*. 2016;22(4):793-801

# Derazantinib inhibits mouse macrophage CSF1R activity

- Derazantinib treatment reduced CSF1-stimulated CSF1R activation (pCSF1R) in a concentration-dependent manner
- The maximum effect is similar to the specific CSF1R inhibitor BLZ945
- Derazantinib active-concentration is achievable in patients

<sup>1</sup> McSheehy et al. Derazantinib (DZB): A dual FGFR/CSF1R-inhibitor active in PDX-models of urothelial cancer. Mol Cancer Ther. 2019 (18) (12 Supplement) LB-C12


## Inhibition of CSF1R activity



*Method: bone-marrow-derived mouse macrophages were starved overnight, treated with CSF1 for 3 min, with or without pre-incubation with BLZ945 or DZB, and then extracted for subsequent immunoblot. The graph shows the quantification of the experiment based on densitometric analysis of the immunoblots*

# Derazantinib — Significant potential beyond iCCA

Frequency of currently known FGFR aberrations across tumor types



Source: Helsten et al., Clin Cancer Res. 2016;22:259-67




# EB1 — A potential response-predictive clinical biomarker for lisavanbulin

- EB1 (plus-end binding protein)<sup>1</sup> is located on the microtubules and involved in microtubule dynamics
- Predictive of response to lisavanbulin in mouse models<sup>1</sup>


<sup>1</sup> Berges et al. EB1-dependent long survival of glioblastoma cancer stem-like cell tumor-bearing mice after oral treatment with the novel tubulin-binding checkpoint activator BAL101553. Eur. J. Cancer 2018, 103, E61-62, A166

## Effect of lisavanbulin (BAL101553) on survival in mice with EB1-expressing or EB1 downregulated GBM

### EB1-expressing GBM




### EB1-downregulated GBM




# Extension of Pfizer supply period

- Supply API and bulk Cresemba vials 2020/2021
    - Increase in product sales (in CHF)
    - Increase in cost of products sold (in CHF); economies-of-scale in supply to other partners
    - Lower gross margin (in % of product sales)
    - Temporary increase in working capital
- => Net positive cash flow over 2020/2021



# Convertible bond transactions — successfully improved debt maturity profile (in CHF mn)



# Glossary

- ABSSSI: **A**cute **b**acterial **s**kin and **s**kin **s**tructure **i**nfections
- CSF1R: **C**olony-**s**timulating **f**actor **1** **r**eceptor
- EAP: **E**xpanded **a**ccess **p**rogram
- FGFR: **F**ibroblast **g**rowth **f**actor **r**eceptor
- FIDES: **F**ibroblast growth factor inhibition with **d**erazantinib in **s**olid tumors
- iCCA: **I**ntrahepatic **c**holangiocarcinoma
- MSSA: **M**ethicillin-**s**usceptible ***S**ta**h**phylococcus **a**ureus*
- MRSA: **M**ethicillin-**r**esistant ***S**ta**h**phylococcus **a**ureus*
- ORR: **O**bjective **r**esponse **r**ate
- PFS: **P**rogression-**f**ree **s**urvival
- SAB: ***S**ta**h**phylococcus **a**ureus* **b**acteremia
- VEGFR2: **V**ascular **e**ndothelial **g**rowth **f**actor **r**eceptor **2**

# Disclaimer and forward-looking statements

This communication, including the accompanying oral presentation, contains certain forward-looking statements, including, without limitation, statements containing the words “believes”, “anticipates”, “expects”, “supposes”, “considers”, and words of similar import, or which can be identified as discussions of strategy, plans or intentions. Such forward-looking statements are based on the current expectations and belief of company management, and are subject to numerous risks and uncertainties, which may cause the actual results, financial condition, performance, or achievements of Basilea, or the industry, to be materially different from any future results, performance, or achievements expressed or implied by such forward-looking statements. Such factors include, among others, the following: the uncertainty of pre-clinical and clinical trials of potential products, limited supplies, future capital needs and the uncertainty of additional funding, compliance with ongoing regulatory obligations and the need for regulatory approval of the company’s operations and potential products, dependence on licenses, patents, and proprietary technology as well as key suppliers and other third parties, including in preclinical and clinical trials, acceptance of Basilea’s products by the market in the event that they obtain regulatory approval, competition from other biotechnology, chemical, and pharmaceutical companies, attraction and retention of skilled employees and dependence on key personnel, and dependence on partners for commercialization of products, limited manufacturing resources, management’s discretion as to the use of proceeds, risks of product liability and limitations on insurance, uncertainties relating to public health care policies, adverse changes in governmental rules and fiscal policies, changes in foreign currency and other factors referenced in this communication. Given these uncertainties, prospective investors are cautioned not to place undue reliance on such forward-looking statements. Basilea disclaims any obligation to update any such forward-looking statements to reflect future events or developments, except as required by applicable law. Derazantinib and its uses are investigational and have not been approved by a regulatory authority for any use. Efficacy and safety have not been established. The information presented should not be construed as a recommendation for use. The relevance of findings in nonclinical/preclinical studies to humans is currently being evaluated.



**Focused on Growth and Innovation**

**Grenzacherstrasse 487  
PO Box  
4005 Basel  
Switzerland**

**investor\_relations@basilea.com  
www.basilea.com**

All rights reserved.  
© Basilea Pharmaceutica International Ltd. 2021